

Supplemental Material

A DNA-DNA hybridisation study of strains of the Flavobacteriaceae family from mostly human clinical sources and proposals of *Chryseobacterium bernardetii* sp. nov., *Chryseobacterium carnis* sp. nov., *Chryseobacterium lactis* sp. nov., *Chryseobacterium nakagawai* sp. nov., and *Chryseobacterium taklimakanense* comb. nov.

By B. Holmes,¹ A. G. Steigerwalt² and A. C. Nicholson²

¹National Collection of Type Cultures, Health Protection Agency, Colindale, London NW9 5EQ, UK

²Special Bacteriology Reference Laboratory, Centers for Disease Control and Prevention, Atlanta, Georgia 30333, USA

Table S1: Sequences of primers used for sequencing the 16S rRNA gene.

Name	Sequence
fD1-5p	CCGAATTCTCGTCGACAACAG
fD1	AGAGTTGATCMTGGCTCAG
F785	GGATTAGATACCCTGGTA
R357	CTGCTGCCTCCCGTA
R802	TACCAGGGTATCTAATCC
F357	TACGGGAGGCAGCAG
R530	GTATTACCGCGGCTGCTG
R519	GWATTACCGCGGCKGCTG
F530	CAGCAGCCCGCGTAATAC
BSF917	GAATTGACGGGRCCC
BSR926	CCGTCATTYYTTTRAGTTT
BSF1099	GYAACGAGCGCAACCC
BSR1114	GGTTGCGCTCGTTRC
BSF1391	TGTACACACCGCCCCGTC
BSR1407	GACGGGCGGTGTGTRC
rP2-5p	CCCGGGATCCAAGCTTAC

Figure S1: A rooted Neighbour-joining phylogenetic tree showing the relationships between type strains of all *Chryseobacterium* species using 1336 bp of 16S rRNA gene sequence with *Weeksella virosa* as the outgroup.

